## Record Display Form

First Hit

Previous Doc

Next Doc Go to Doc#

Generate Collection

L15: Entry 52 of 59

File: DWPI

Jan 16, 1996

DERWENT-ACC-NO: 1996-112624

DERWENT-WEEK: 199612

COPYRIGHT 2005 DERWENT INFORMATION LTD

TITLE: Dermal external preparation having melanin formation-inhibitory action -

contains extract of Achote, useful for treating spots, freckles, etc.

PATENT-ASSIGNEE:

**ASSIGNEE** 

CODE

SHISEIDO CO LTD

SHIS

PRIORITY-DATA: 1994JP-0168689 (June 29, 1994)

Search Selected

Search ALL

Clear

PATENT-FAMILY:

PUB-NO

PUB-DATE

LANGUAGE

PAGES

MAIN-IPC

JP 08012563 A

January 16, 1996

007

A61K007/48

APPLICATION-DATA:

PUB-NO

APPL-DATE

APPL-NO

DESCRIPTOR

JP 08012563A

June 29, 1994

1994JP-0168689

INT-CL (IPC): A61 K  $\frac{7}{00}$ ; A61 K  $\frac{7}{48}$ ; A61 K  $\frac{35}{78}$ 

ABSTRACTED-PUB-NO: JP 08012563A

BASIC-ABSTRACT:

Dermal external prepn. contains extract of Achote, Bixa orellana L., a South American xeromorphic grassland plant, partic. at ratios of 0.005-20.0 wt.% and is used for skin melanocyte inhibitor.

Extract of Achote, Bixa orellana L., with organic solvent (e.g. MeOH, EtOH, aq. alcohols, acetone and EtOAc) is added to cosmetics base at concn. of 0.005-20.0 (pref. 0.01-10.0) dried wt.% together with other conventional additives (e.g. humectants, antioxidants, UV-absorption agents, thickener and skin nutrients). It is prepd. in conventional forms including ointments, cream, lotion, pack and bathing agents.

USE/ADVANTAGE - Used to treat and prevent spots, freckles, chloasma and sunburn. The treatment agent has melanin formation and tyrosinase inhibitory activity.

In an example, mouse derived B16 melanoma cells were cultured in 95% O2 and 5% CO2 atmos. at 37 deg. C. Extract was added at concns. of 1/100-100000 wt.%, cultured for further 3 days and microscopically observed. Slight inhibition of melanin formation was observed at a concn. of 1/100000 wt.% and tyrosinase inhibitory